

AN S_3 -SYMMETRIC LITTLEWOOD-RICHARDSON RULE

HUGH THOMAS AND ALEXANDER YONG

1. INTRODUCTION

Fix Young shapes $\lambda, \mu, \nu \subseteq \Lambda := \ell \times k$. Viewing the **Littlewood-Richardson coefficients** $C_{\lambda, \mu, \nu}$ as intersection numbers of three Schubert varieties in general position in the Grassmannian of ℓ -dimensional planes in $\mathbb{C}^{\ell+k}$, one has the obvious S_3 -symmetries:

$$(1) \quad C_{\lambda, \mu, \nu} = C_{\mu, \nu, \lambda} = C_{\nu, \lambda, \mu} = C_{\mu, \lambda, \nu} = C_{\nu, \mu, \lambda} = C_{\lambda, \nu, \mu}.$$

There is interest in the combinatorics of these symmetries; previously studied Littlewood-Richardson rules for $C_{\lambda, \mu, \nu}$ manifest at most three of the six, see, e.g., [BeZe91, KnTaWo01, VaPa05, HeKa06] and the references therein. We construct a **carton rule** for $C_{\lambda, \mu, \nu}$ that transparently and uniformly explains all symmetries (1).

Figure 1 depicts a **carton**, i.e., a three-dimensional box with a grid drawn rectilinearly on the six faces of its surface. Along the “ $\emptyset, T_\lambda, T_\mu$ ” face, the grid has $(|\mu| + 1) \times (|\lambda| + 1)$ vertices (including ones on the bounding edges of the face); the remainder of the grid is similarly determined. Define a **carton filling** to be an assignment of a Young diagram to each vertex of the grid so the shapes increase one box at a time while moving away from

\emptyset , and so that, for any subgrid $\begin{array}{c} \alpha \quad - \quad \beta \\ | \quad \quad | \\ \gamma \quad - \quad \delta \end{array}$ the **Fomin growth assumptions** hold:

- (F1) if α is the unique shape containing γ and contained in β , then $\delta = \alpha$;
- (F2) otherwise there is a unique such shape other than α , and this shape is δ .

Notice that these conditions are symmetric in α and δ .

Initially, assign the shapes \emptyset and Λ to opposite corners, as in Figure 1. A standard Young tableau $T \in \text{SYT}(\sigma/\pi)$ of shape σ/π is equivalent to a **shape chain** in Young’s lattice, e.g.,

$$T = \begin{array}{|c|c|c|c|} \hline & & & 1 \ 5 \\ \hline & 2 \ 3 & & \\ \hline 4 & & & \\ \hline \end{array} \leftrightarrow \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array} - \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} - \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} - \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} - \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline \end{array} - \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline \end{array}$$

by starting with the unfilled inner shape, and adding one box at a time to indicate the box containing 1, then 2, etc. Fix a choice of standard tableaux T_λ, T_μ and T_ν of respective shapes λ, μ and ν . Initialize the indicated edges with the shape chains for these tableaux.

Let $\text{CARTONS}_{\lambda, \mu, \nu}$ be all carton fillings with the above initial data.

Main Theorem. *The Littlewood-Richardson coefficient $C_{\lambda, \mu, \nu}$ equals $\#\text{CARTONS}_{\lambda, \mu, \nu}$.*

This rule manifests bijections between $\text{CARTONS}_{\lambda, \mu, \nu}$ and $\text{CARTONS}_{\delta, \epsilon, \zeta}$ for permutations $(\delta, \epsilon, \zeta)$ of (λ, μ, ν) . We remark that it can be stated as counting lattice points of a 0, 1-polytope, and readily extends to the setting of [ThYo06, ThYo07].

Date: April 4, 2007.




FIGURE 1. The carton rule counts $C_{\lambda,\mu,\nu}$ by assigning Young diagrams to the vertices of the six faces

The Theorem is proved in Section 2, starting from the jeu de taquin formulation of the Littlewood-Richardson rule, and using Fomin's growth diagram ideas. In Figure 2 we give an example of the Main Theorem. An extended example is given in Section 3.




FIGURE 2. The “front” three faces of a carton filling for $C_{(2),(2,1),(1)} = 1$. The choices of tableaux T_λ , T_μ and T_ν are as shown.

2. TABLEAU FACTS AND PROOF OF THE MAIN THEOREM

2.1. Tableau sliding. There is a partial order \prec on the rectangle Λ where $x \prec y$ if x is weakly northwest of y . Given $T \in \text{SYT}(\nu/\lambda)$ consider $x \in \lambda$, maximal in \prec subject to being less than *some* box of ν/λ . Associate another standard tableau $\text{jdt}_x(T)$, called the **jeu de taquin slide** of T into x : Let y be the box of ν/λ with the smallest label, among those covering x . Move the label of y to x , leaving y vacant. Look for boxes of ν/λ covering y and repeat, moving into y the smallest label among those boxes covering it. Then $\text{jdt}_x(T)$ results when no further slides are possible. The **rectification** of T is the iteration of jeu de taquin slides until terminating at a straight shape standard tableau $\text{rectification}(T)$.

We can impose a specific “inner” $U \in \text{SYT}(\lambda)$ that encodes the order in which the jeu de taquin slides are done. For example, if $U = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & & \\ \hline \end{array}$ then

$$T = \begin{array}{|c|c|c|c|c|} \hline & & & 1 & 5 \\ \hline & 2 & 3 & & \\ \hline 4 & 6 & & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|c|} \hline & & & 1 & 5 \\ \hline 2 & 3 & & & \\ \hline 4 & 6 & & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|c|} \hline & & 1 & 5 & \\ \hline 2 & 3 & & & \\ \hline 4 & 6 & & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|c|} \hline & 1 & 5 & & \\ \hline 2 & 3 & & & \\ \hline 4 & 6 & & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|c|} \hline 1 & 3 & 5 & & \\ \hline 2 & 6 & & & \\ \hline 4 & & & & \\ \hline \end{array}$$

$= \text{rectification}(T)$. Placing these chains one atop another (starting with T 's and ending with $\text{rectification}(T)$'s) gives a Fomin **growth diagram**, which in this case is given in Table 1. Growth diagrams satisfy (F1) and (F2); see Fomin's [St99, Appendix 2] for more.

(3, 1)	(4, 1)	(4, 2)	(4, 3)	(4, 3, 1)	(5, 3, 1)	(5, 3, 2)
(3)	(4)	(4, 1)	(4, 2)	(4, 2, 1)	(5, 2, 1)	(5, 2, 2)
(2)	(3)	(3, 1)	(3, 2)	(3, 2, 1)	(4, 2, 1)	(4, 2, 2)
(1)	(2)	(2, 1)	(2, 2)	(2, 2, 1)	(3, 2, 1)	(3, 2, 2)
\emptyset	(1)	(1, 1)	(2, 1)	(2, 1, 1)	(3, 1, 1)	(3, 2, 1)

TABLE 1. A growth diagram

Given a top row U and left column T , define $\text{infusion}_1(U, T)$ to be the bottom row of the growth diagram, and $\text{infusion}_2(U, T)$ to be the right column. We write $\text{infusion}(U, T)$ for the ordered pair $(\text{infusion}_1(U, T), \text{infusion}_2(U, T))$. Growth diagrams are transpose symmetric, because (F1) and (F2) are. So one has the **infusion involution**:

$$\text{infusion}(\text{infusion}(U, T)) = (U, T).$$

If U is a straight shape, $\text{infusion}_1(U, T) = \text{rectification}(T)$, while $\text{infusion}_2(U, T)$ encodes the order in which squares were vacated in the jeu de taquin process.

Also, given $T \in \text{SYT}(\nu/\lambda)$, consider $x \in \Lambda \setminus \nu$ *minimal* subject to being larger than some element of ν/λ . The **reverse jeu de taquin slide** $\text{revjdt}_x(T)$ of T into x is defined similarly to a jeu de taquin slide, except we move into x the *largest* of the labels among boxes in ν/λ covered by x . We define **reverse rectification** $\text{revrectification}(T)$ similarly. The first fundamental theorem of jeu de taquin asserts $(\text{rev})\text{rectification}$ is well-defined.

Fix $T_\mu \in \text{SYT}(\mu)$. The number of $T \in \text{SYT}(\nu/\lambda)$ such that $\text{rectification}(T) = T_\mu$ equals $C_{\lambda, \mu, \nu^\vee} = C_{\lambda, \mu}^\vee$ where ν^\vee is the straight shape obtained as the 180-degree rotation of $\Lambda \setminus \nu$. This is Schützenberger's jeu de taquin formulation of the Littlewood-Richardson rule, see [St99, Appendix 2].

By a **slide** we mean either kind of jeu de taquin slide. Consider two equivalence relations on a pair of tableaux T and U . Tableaux are **jeu de taquin equivalent** if one can be

obtained from the other by a sequence of slides. They are **dual equivalent** if any such sequence results in tableaux of the same shape [Ha92]. Facts: Tableaux of the same straight shape are dual equivalent. A common application of slides to dual equivalent tableaux produces dual equivalent tableaux. A pair of tableaux that are both jeu de taquin and dual equivalent must be equal.

Recall Schützenberger’s evacuation map. For $T \in \text{SYT}(\lambda)$, let \hat{T} be obtained by erasing the entry 1 (in the northwest corner c) of T and subtracting 1 from the remaining entries. Let $\Delta(T) = \text{jdt}_c(\hat{T})$. The **evacuation** $\text{evac}(T) \in \text{SYT}(\lambda)$ is defined by the shape chain

$$\emptyset = \text{shape}(\Delta^{|\lambda|}(T)) - \text{shape}(\Delta^{|\lambda|-1}(T)) - \dots - \text{shape}(\Delta^1(T)) - T.$$

This map is an involution: $\text{evac}(\text{evac}(T)) = T$.

2.2. Proof of the rule. Given $T \in \text{SYT}(\alpha)$ for a straight shape α , define $\tilde{T} \in \text{SYT}(\text{rotate}(\alpha))$ where $\text{rotate}(\alpha) = \Lambda \setminus \alpha^\vee$ by computing $\text{evac}(T) \in \text{SYT}(\alpha)$, replacing entry i with $|\alpha| - i + 1$ throughout and rotating the resulting tableau 180-degrees and placing it at the bottom right corner of Λ .

We need the following well-known fact. The proof we give extends straightforwardly to the setup of [ThYo06, ThYo07], allowing a “cominuscule” version of the Main Theorem.

Lemma 2.1. *Suppose α, β and γ are shapes where*

$$T_\beta \in \text{SYT}(\beta), T_{\gamma^\vee/\alpha} \in \text{SYT}(\gamma^\vee/\alpha) \text{ and } \text{rectification}(T_{\gamma^\vee/\alpha}) = T_\beta.$$

Then $\text{revrectification}(T_{\gamma^\vee/\alpha}) = \text{revrectification}(T_\beta) = \tilde{T}_\beta$.

Proof. It suffices to show that $\text{revrectification}(T_\beta) = \tilde{T}_\beta$. We induct on $|\beta| = n$.

Let U be the tableau obtained by removing the box labeled 1 from T_β . So U is of skew shape $\beta/(1)$. Let $V = \text{rectification}(U)$, a tableau of shape $\kappa \subset \beta$. By induction, $\text{revrectification}(V) = \tilde{V}$. Now $\text{revrectification}(T_\beta)$ and $\text{revrectification}(V)$ agree except that the former has a label 1 in a box that does not appear in the latter. In [ThYo06, Proposition 4.6], we showed that the reverse rectification of a tableau of shape β is necessarily of shape $\text{rotate}(\beta)$. Thus, the location of 1 in $\text{revrectification}(T_\beta)$ must be the box κ^\vee/β^\vee . This is the 180-degree rotation of the box β/κ in Λ , the latter being the position of n in $\text{evac}(T_\beta)$. Thus, 1 is located in the desired position in $\text{revrectification}(T_\beta)$. The rest of the statement follows from the fact that the rest of $\text{revrectification}(T_\beta)$ agrees with \tilde{V} , and the entries of $\text{evac}(T_\beta)$ other than n agree with $\text{evac}(V)$. \square

Corollary 2.2. *Fix a carton filling. The uninitialized corners of the “ $\emptyset, T_\lambda, T_\mu$ ” and “ $\emptyset, T_\nu, T_\lambda$ ” faces are ν^\vee and μ^\vee respectively. The remaining “sixth” corner (the unique one not visible in Figure 1) is λ^\vee . Thus, we can speak of the edges $\lambda^\vee - \Lambda$, $\mu^\vee - \Lambda$ and $\nu^\vee - \Lambda$. These represent the shape chains of \tilde{T}_λ , \tilde{T}_μ and \tilde{T}_ν respectively.*

Proof. Think of the union of the faces “ $\emptyset, T_\mu, T_\lambda$ ” with the adjacent face involving Λ and the “sixth corner” as a single growth diagram. This diagram computes $\text{revrectification}(T_\lambda)$ and records the result along the edge involving Λ and diagonally opposite to the T_λ edge. But the lemma asserts this result is \tilde{T}_λ and hence the “sixth corner vertex” is assigned λ^\vee . The other conclusions are proved similarly. \square

Corollary 2.2 affords us the convenience of referring to a face by its corner vertices. Note any carton filling gives a growth diagram on the face $\emptyset - \mu - \nu^\vee - \lambda$ for which the edge $\lambda - \nu^\vee$ is a standard tableau of shape ν^\vee/λ rectifying to T_μ . By the jeu de taquin Littlewood-Richardson rule, fillings of this face count $C_{\lambda,\mu,\nu}$. Hence it suffices to show that any such growth diagram for this face extends uniquely to a filling of the entire carton.

If such an extension exists, it is unique: $\lambda - \nu^\vee$ and $\Lambda - \nu^\vee$ determine, by (F1), (F2) and the Corollary, the face $\lambda - \mu^\vee - \Lambda - \nu^\vee$. Similarly, these two faces determine the remaining faces (in order):

$$\emptyset - \nu - \mu^\vee - \lambda \implies \nu - \lambda^\vee - \Lambda - \mu^\vee \implies \mu - \lambda^\vee - \Lambda - \nu^\vee \implies \emptyset - \nu - \lambda^\vee - \mu.$$

We now show that with a filling of $\emptyset - \mu - \nu^\vee - \lambda$ (together with the extra edges given in the Corollary) one can extend it to fill the carton, using the conclusions of Corollary 2.2.

$\emptyset - \mu - \nu^\vee - \lambda$: This is given by labeling $\lambda - \nu^\vee$ with $T_{\nu^\vee/\lambda} \in \text{SYT}(\nu^\vee/\lambda)$ that **witnesses** $C_{\lambda,\mu,\nu} = C_{\lambda,\mu}^{\nu^\vee}$ i.e., it rectifies to T_μ . By the infusion involution, the edge $\mu - \nu^\vee$ representing $T_{\nu^\vee/\mu} := \text{infusion}_2(T_\lambda, T_{\nu^\vee/\lambda}) \in \text{SYT}(\nu^\vee/\mu)$ witnesses $C_{\mu,\lambda}^{\nu^\vee}$.

$\lambda - \mu^\vee - \Lambda - \nu^\vee$: Build the growth diagram using $T_{\nu^\vee/\lambda}$ and \tilde{T}_ν from the edges $\lambda - \nu^\vee$ and $\nu^\vee - \Lambda$ respectively. The only boundary condition we need to check is that $\text{infusion}_2(T_{\nu^\vee/\lambda}, \tilde{T}_\nu) = \tilde{T}_\mu$, which is true by the Lemma. The newly determined edge $\lambda - \mu^\vee$ is $\text{infusion}_1(T_{\nu^\vee/\lambda}, \tilde{T}_\nu)$. This is a tableau $T_{\mu^\vee/\lambda} \in \text{SYT}(\mu^\vee/\lambda)$ which rectifies to T_ν , i.e., one that witnesses $C_{\lambda,\nu}^{\mu^\vee}$.

$\emptyset - \nu - \mu^\vee - \lambda$: Using the determined edges $\lambda - \mu^\vee$ and $\emptyset - \lambda$ we obtain a growth diagram with edge $\emptyset - \nu$ representing $\text{infusion}_1(T_\lambda, T_{\mu^\vee/\lambda})$, which equals T_ν , as desired. By the infusion involution, $\nu - \mu^\vee$ represents a tableau $T_{\mu^\vee/\nu} \in \text{SYT}(\mu^\vee/\nu)$ that witnesses $C_{\nu,\lambda}^{\mu^\vee}$.

$\nu - \lambda^\vee - \Lambda - \mu^\vee$: This time, we grow the face using the edges $\nu - \mu^\vee$ and $\mu^\vee - \Lambda$. The edge $\lambda^\vee - \Lambda$ is thus $\text{infusion}_2(T_{\mu^\vee/\nu}, \tilde{T}_\mu)$ which indeed equals \tilde{T}_λ , by the Lemma. The newly determined edge $\nu - \lambda^\vee$ is $T_{\lambda^\vee/\nu} := \text{infusion}_1(T_{\mu^\vee/\nu}, \tilde{T}_\mu) \in \text{SYT}(\lambda^\vee/\nu)$ that witnesses $C_{\nu,\mu}^{\lambda^\vee}$.

$\mu - \lambda^\vee - \Lambda - \nu^\vee$: Growing the face using $\mu - \nu^\vee$ and $\nu^\vee - \Lambda$ we find that the edge $\lambda^\vee - \Lambda$ equals $\text{infusion}_2(T_{\nu^\vee/\mu}, \tilde{T}_\nu)$, which is the already determined \tilde{T}_λ . The newly determined edge $\mu - \lambda^\vee$ is $T_{\lambda^\vee/\mu} := \text{infusion}_1(T_{\nu^\vee/\mu}, \tilde{T}_\nu) \in \text{SYT}(\lambda^\vee/\mu)$ that witnesses $C_{\mu,\nu}^{\lambda^\vee}$.

$\emptyset - \nu - \lambda^\vee - \mu$: We grow this final face using $\emptyset - \nu$ and $\nu - \lambda^\vee$, but need to make two consistency checks. First the edge $\emptyset - \mu$ is $\text{infusion}_1(T_\nu, T_{\lambda^\vee/\nu})$ which clearly is T_μ .

It remains to check that

$$(2) \quad \text{infusion}_2(T_\nu, T_{\lambda^\vee/\nu}) = T_{\lambda^\vee/\mu}.$$

(Notice that $\text{infusion}_2(T_\nu, T_{\lambda^\vee/\nu})$ is a filling of λ^\vee/μ that rectifies to T_ν , just as $T_{\lambda^\vee/\mu}$ does. However it is not clear *a priori* that they are the same.)

To prove (2), we need a definition. For tableaux A and B of respective (skew) shapes α and γ/α let $A \star B$ be their concatenation as a (nonstandard) tableau. If C is a tableau of shape Λ/γ and α is a straight shape, $A \star B \star C$ is a **layered tableau** of shape Λ .

Example 2.3. Let $\alpha = (2, 1)$ and $\gamma = (4, 2, 1)$. We have

$$A = \begin{array}{|c|c|} \hline \mathbf{1} & \mathbf{2} \\ \hline 3 & \\ \hline \end{array}, \quad B = \begin{array}{|c|c|c|c|} \hline & & \mathbf{2} & \mathbf{3} \\ \hline & \mathbf{1} & & \\ \hline 4 & & & \\ \hline \end{array}, \quad C = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & \mathbf{2} & \mathbf{3} \\ \hline & \mathbf{1} & \mathbf{4} & \mathbf{5} \\ \hline \end{array}, \quad \text{and } A \star B \star C = \begin{array}{|c|c|c|c|} \hline \mathbf{1} & \mathbf{2} & \mathbf{2} & \mathbf{3} \\ \hline 3 & \mathbf{1} & \mathbf{2} & \mathbf{3} \\ \hline 4 & \mathbf{1} & \mathbf{4} & \mathbf{5} \\ \hline \end{array}.$$

We have used boldface and underlining to distinguish the entries from A , B and C .

Let \mathcal{I}_1 and \mathcal{I}_2 be operators on layered tableaux defined by

$$\mathcal{I}_1 : A \star B \star C \mapsto \text{infusion}_1(A, B) \star \text{infusion}_2(A, B) \star C$$

and

$$\mathcal{I}_2 : A \star B \star C \mapsto A \star \text{infusion}_1(B, C) \star \text{infusion}_2(B, C).$$

The infusion involution says \mathcal{I}_1^2 and \mathcal{I}_2^2 are the identity operator. In fact, the following crucial “braid identity” holds, showing \mathcal{I}_1 and \mathcal{I}_2 generate a representation of S_3 :

Proposition 2.4. $\mathcal{I}_1 \circ \mathcal{I}_2 \circ \mathcal{I}_1 = \mathcal{I}_2 \circ \mathcal{I}_1 \circ \mathcal{I}_2$.

In view of the Proposition, (2) follows by setting $\alpha = \lambda$, $\gamma = \nu^\vee$, $A = T_\lambda$, $B = T_{\lambda^\vee/\nu}$ and $C = \tilde{T}_\nu$, since the assertion merely says the “middle” tableau in

$$(3) \quad \mathcal{I}_1 \circ \mathcal{I}_2 \circ \mathcal{I}_1 \circ \mathcal{I}_2(T_\lambda \star T_{\nu^\vee/\lambda} \star \tilde{T}_\nu) \text{ and } \mathcal{I}_2 \circ \mathcal{I}_1(T_\lambda \star T_{\nu^\vee/\lambda} \star \tilde{T}_\nu)$$

are the same, whereas we even have equality of the two layered tableaux.

Proof of Proposition 2.4: By the Lemma it follows that

$$(4) \quad \tilde{C} \star \bar{B} \star \tilde{A} := \mathcal{I}_1 \circ \mathcal{I}_2 \circ \mathcal{I}_1(A \star B \star C), \text{ and}$$

$$(5) \quad \tilde{C} \star \hat{B} \star \tilde{A} := \mathcal{I}_2 \circ \mathcal{I}_1 \circ \mathcal{I}_2(A \star B \star C),$$

where the shapes of \tilde{C} , \bar{B} and \tilde{A} are respectively the 180 degree rotations of the shapes of A , B and C , and we know the fillings of \tilde{C} and \tilde{A} in terms of evacuation. We thus also know the shapes of \bar{B} and \hat{B} are the same. It remains to show \bar{B} equals \hat{B} .

We first study (4). Let $B' = \text{infusion}_1(A, B)$, and $A' = \text{infusion}_2(A, B)$. Now

$$(6) \quad \mathcal{I}_1 \circ \mathcal{I}_2(B' \star A' \star C) = \tilde{C} \star \bar{B} \star \tilde{A},$$

and the skew tableau $\bar{B} \star \tilde{A}$ is obtained by treating $B' \star A'$ as a *single standard tableau* by valuing an entry i of A' as $|B'| + i$, evacuating, rotating the result 180 degrees and finally sending the entry i from A' (respectively B') to $|A'| - i + 1$ (respectively $|B'| - i + 1$).

Example 2.5. Continuing the previous example,

$$B' \star A' = \begin{array}{|c|c|c|c|} \hline \mathbf{1} & \mathbf{2} & \mathbf{3} & \mathbf{2} \\ \hline 4 & 3 & & \\ \hline 1 & & & \\ \hline \end{array} \text{ and } \text{evac}(B' \star A') = A'' \star B'' = \begin{array}{|c|c|c|c|} \hline \mathbf{1} & \mathbf{3} & \mathbf{3} & \mathbf{4} \\ \hline 2 & \mathbf{1} & & \\ \hline 2 & & & \\ \hline \end{array} \mapsto \bar{B} \star \tilde{A} = \begin{array}{|c|c|c|c|} \hline & & & \mathbf{3} \\ \hline & & 4 & 2 \\ \hline 1 & 2 & 1 & 3 \\ \hline \end{array}$$

Let us focus on $\text{evac}(B' \star A') = A'' \star B''$ and momentarily ignore the subsequent rotation and complementation of entries. Fomin has shown in [St99, Appendix 2] that a fruitful way to think of $\text{evac}(X)$ is with triangular growth diagrams obtained by placing the shape chain of X , then $\Delta(X)$, $\Delta^2(X)$, \dots on top of one another (slanted left to right); see Figure 3.

We begin with the data $B' \star A'$ along the left side of the diagram. This is given as a concatenation of two shape chains, one from $\emptyset - \mu$ and then one from $\mu - \gamma$ where μ is the shape of B' . Applying (F1) and (F2) from Section 1, we obtain $A'' \star B'' = \text{evac}(B' \star A')$,




FIGURE 3. A triangular growth diagram to compute $\text{evac}(B' \star A')$

given along the righthand side as a pair of chains connected at α . It follows from the construction of growth diagrams that the thick lines represent $\text{evac}(B')$ and $A = \text{evac}(A'')$ (the latter being a consequence of the Lemma).

These thick lines together with the vertex γ define a rectangular growth diagram, so:

$$(7) \quad \text{infusion}(A, B'') = (\text{evac}(B'), A').$$

In particular, the shape of $\text{rectification}(B'') = \mu$. On the other hand, by definition

$$(8) \quad \text{infusion}(A, B) = (B', A'),$$

and B rectifies to a tableau of shape μ also.

By (7) and (8) combined with the aforementioned results of [Ha92] we see B and B'' are dual equivalent, as they both are obtained by an application of the same sequence of slides (encoded by A') to a pair of tableaux of the same straight shape. Moreover, by (7) B'' is jeu de taquin equivalent to $\text{evac}(B') = \text{evac}(\text{rectification}(B))$. Carrying out the analogous “reverse” analysis on $\mathcal{I}_2 \circ \mathcal{I}_1 \circ \mathcal{I}_2(A \star B \star C)$ we let $C^\circ = \text{infusion}_1(B, C)$ and $B^\circ = \text{infusion}_2(B, C)$ and study $\mathcal{I}_2 \circ \mathcal{I}_1(A \star C^\circ \star B^\circ) = \tilde{C} \star \hat{B} \star \tilde{A}$. Parallel to (6) we consider the “reverse evacuation” $B^{\circ\circ} \star C^{\circ\circ}$ of $C^\circ \star B^\circ$ using *reverse* jeu de taquin slides. Then we similarly conclude $B^{\circ\circ}$ is dual equivalent to B (and thus B''). Also $B^{\circ\circ}$ is jeu de taquin equivalent to the reverse evacuation of $\text{revrectification}(B)$. Thus by the lemma, B'' and $B^{\circ\circ}$ are jeu de taquin equivalent. Hence $B'' = B^{\circ\circ}$ and so $\bar{B} = \hat{B}$, as required.

Example 2.6. Revisiting Example 2.3, $C^\circ \star B^\circ =$

	2	3	
4	5	3	
1	1	2	4

 with reverse evacuation

3	4		
1	1	2	
2	3	4	5

 and $B^{\circ\circ}$ (boldface in the latter tableau) is also B'' from Example 2.5. Rotation and complementation gives \hat{B} , which is \bar{B} from Example 2.5, as desired. \square

3. AN EXTENDED EXAMPLE OF THE MAIN THEOREM

Let $\Lambda = 3 \times 4 =$

, $\lambda = (2, 1) =$

, $\mu = (3, 1) =$

, $\nu = (3, 2) =$

.

Therefore $\lambda^\vee = (4, 3, 2)$, $\mu^\vee = (4, 3, 1)$ and $\nu^\vee = (4, 2, 1)$. Also

$$T_\lambda = \begin{bmatrix} 1 & 2 \\ 3 \end{bmatrix}, T_\mu = \begin{bmatrix} 1 & 2 & 3 \\ 4 \end{bmatrix}, T_\nu = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 \end{bmatrix}, \tilde{T}_\lambda = \begin{bmatrix} & & & 1 \\ & & & 2 \\ & & 2 & 3 \end{bmatrix}, \tilde{T}_\mu = \begin{bmatrix} & & & 3 \\ & & & 1 \\ & 1 & 2 & 4 \end{bmatrix}, \text{ and } \tilde{T}_\nu = \begin{bmatrix} & & & \\ & & 2 & 3 \\ & 1 & 4 & 5 \end{bmatrix}.$$

Here $C_{\lambda, \mu, \nu} = 1$, and we now give the unique carton filling. We begin with $T_{\nu^\vee/\lambda} =$

		2	3
	4		
1			

Then we have the following sides of the carton, described as growth diagrams with the obvious identifications of boundaries (the partitions correspond to shapes placed on the vertices of the carton and the diagrams have been oriented to be consistent with Figure 1):

$(2, 1) = \lambda$	$(2, 1, 1)$	$(3, 1, 1)$	$(4, 1, 1)$	$(4, 2, 1) = \nu^\vee$
(2)	$(2, 1)$	$(3, 1)$	$(4, 1)$	$(4, 2)$
(1)	$(1, 1)$	$(2, 1)$	$(3, 1)$	$(3, 2)$
\emptyset	(1)	(2)	(3)	$(3, 1) = \mu$

TABLE 2. $\emptyset - \mu - \nu^\vee - \lambda$

$(4, 3, 1) = \mu^\vee$	$(4, 3, 2)$	$(4, 3, 3)$	$(4, 4, 3)$	$(4, 4, 4) = \Lambda$
$(4, 2, 1)$	$(4, 2, 2)$	$(4, 3, 2)$	$(4, 4, 2)$	$(4, 4, 3)$
$(4, 2)$	$(4, 2, 1)$	$(4, 3, 1)$	$(4, 4, 1)$	$(4, 4, 2)$
$(3, 2)$	$(3, 2, 1)$	$(3, 3, 1)$	$(4, 3, 1)$	$(4, 3, 2)$
$(2, 2)$	$(2, 2, 1)$	$(3, 2, 1)$	$(4, 2, 1)$	$(4, 2, 2)$
$(2, 1) = \lambda$	$(2, 1, 1)$	$(3, 1, 1)$	$(4, 1, 1)$	$(4, 2, 1) = \nu^\vee$

TABLE 3. $\lambda - \mu^\vee - \Lambda - \nu^\vee$

$(4, 3, 1) = \mu^\vee$	$(4, 2, 1)$	$(4, 2)$	$(3, 2)$	$(2, 2)$	$(2, 1) = \lambda$
$(4, 2, 1)$	$(4, 1, 1)$	$(4, 1)$	$(3, 1)$	$(2, 1)$	(2)
$(3, 2, 1)$	$(3, 1, 1)$	$(3, 1)$	$(2, 1)$	$(1, 1)$	(1)
$(3, 2) = \nu$	$(3, 1)$	(3)	(2)	(1)	\emptyset

TABLE 4. $\emptyset - \nu - \mu^\vee - \lambda$

$(4, 3, 1) = \mu^\vee$	$(4, 3, 2)$	$(4, 3, 3)$	$(4, 4, 3)$	$(4, 4, 4) = \Lambda$
$(4, 2, 1)$	$(4, 2, 2)$	$(4, 3, 2)$	$(4, 4, 2)$	$(4, 4, 3)$
$(3, 2, 1)$	$(3, 2, 2)$	$(3, 3, 2)$	$(4, 3, 2)$	$(4, 4, 2)$
$(3, 2) = \nu$	$(3, 2, 1)$	$(3, 3, 1)$	$(4, 3, 1)$	$(4, 3, 2) = \lambda^\vee$

TABLE 5. $\nu - \lambda^\vee - \Lambda - \mu^\vee$

ACKNOWLEDGMENTS

HT was partially supported by an NSERC Discovery Grant and AY by NSF grant 0601010. This work was partially completed while HT was a visitor at the Centre de Recherches Mathématiques and the University of Minnesota, and while AY was at the Fields Institute and the University of Michigan, Ann Arbor. We thank the Banff International Research Station for a common visit during the “Schubert calculus and Schubert

$(4, 4, 4) = \Lambda$	$(4, 4, 3)$	$(4, 4, 2)$	$(4, 3, 2)$	$(4, 2, 2)$	$(4, 2, 1) = \nu^\vee$
$(4, 4, 3)$	$(4, 4, 2)$	$(4, 4, 1)$	$(4, 3, 1)$	$(4, 2, 1)$	$(4, 2)$
$(4, 3, 3)$	$(4, 3, 2)$	$(4, 3, 1)$	$(3, 3, 1)$	$(3, 2, 1)$	$(3, 2)$
$(4, 3, 2) = \lambda^\vee$	$(4, 2, 2)$	$(4, 2, 1)$	$(3, 2, 1)$	$(3, 1, 1)$	$(3, 1) = \mu$

TABLE 6. $\lambda^\vee - \Lambda - \nu^\vee - \mu$

$(3, 2) = \nu$	$(3, 2, 1)$	$(3, 3, 1)$	$(4, 3, 1)$	$(4, 3, 2) = \lambda^\vee$
$(3, 1)$	$(3, 1, 1)$	$(3, 2, 1)$	$(4, 2, 1)$	$(4, 2, 2)$
(3)	$(3, 1)$	$(3, 2)$	$(4, 2)$	$(4, 2, 1)$
(2)	$(2, 1)$	$(2, 2)$	$(3, 2)$	$(3, 2, 1)$
(1)	$(1, 1)$	$(2, 1)$	$(3, 1)$	$(3, 1, 1)$
\emptyset	(1)	(2)	(3)	$(3, 1) = \mu$

TABLE 7. $\emptyset - \nu - \lambda^\vee - \mu$

geometry” workshop organized by Jim Carrell and Frank Sottile. We also thank Sergey Fomin, Allen Knutson, Ezra Miller, Vic Reiner, Luis Serrano, David Speyer and Dennis Stanton for helpful discussions.

REFERENCES

- [BeZe91] A. D. Berenstein and A. V. Zelevinsky, *Triple multiplicities of $sl(r+1)$ and the spectrum of the exterior algebra of the adjoint representation*, Journal of Algebraic Combinatorics, **1** (1992), 7–22.
- [Ha92] M. Haiman, *Dual equivalence with applications, including a conjecture of Proctor*, Discrete Math. **99** (1992), 79–113.
- [HeKa06] A. Henriques and J. Kamnitzer, *The octahedron recurrence and gl_n crystals*, Adv. Math., **206** (2006), no. 1, 211–249.
- [KnTaWo01] A. Knutson, T. Tao and C. Woodward, *The honeycomb model of $GL(n)$ tensor products II: Puzzles determine facets of the Littlewood-Richardson cone*, J. Amer. Math. Soc., **17** (2004), 19–48.
- [St99] R. P. Stanley, *Enumerative Combinatorics, Volume 2* (with an appendix by S. Fomin), Cambridge University Press, 1999.
- [VaPa05] E. Vallejo and I. Pak, *Combinatorics and geometry of Littlewood-Richardson cones*, Europ. J. Combinat., **26**(2005), 995–1008.
- [ThYo06] H. Thomas and A. Yong, *A combinatorial rule for (co)minuscule Schubert calculus*, preprint math.AG/0608273.
- [ThYo07] ———, *Cominuscule tableau combinatorics*, preprint math.CO/0701215.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW BRUNSWICK, FREDERICTON, NEW BRUNSWICK, E3B 5A3, CANADA

E-mail address: hugh@math.unb.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455, USA

E-mail address: ayong@math.umn.edu